

Montana Health Alert Network

DPHHS HAN

ADVISORY

Cover Sheet

DATE

March 29, 2024

SUBJECT

Increase in Invasive Serogroup Y Meningococcal Disease in the United States.

INSTRUCTIONS

DISTRIBUTE to your local HAN contacts. This HAN is intended for general sharing of information.

- Time for Forwarding: **As Soon As Possible**
- Please forward to DPHHS at hhshan@mt.gov
- **Remove this cover sheet before redistributing and replace it with your own**



For LOCAL HEALTH DEPARTMENT reference only
DPHHS Subject Matter Resource for more information regarding this HAN,

contact:

DPHHS CDCP

Epidemiology Section
1-406-444-0273

Immunization Section
1-406-444-5580

For technical issues related to the HAN message contact the Emergency Preparedness Section at 1-406-444-0919

DPHHS HAN Website:
<https://dphhs.mt.gov/publichealth/han>

REMOVE THIS COVER SHEET BEFORE REDISTRIBUTING AND REPLACE IT WITH YOUR OWN

Please ensure that DPHHS is included on your HAN distribution list.
hhshan@mt.gov

Categories of Health Alert Messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.

Information Service: passes along low level priority messages that do not fit other HAN categories and are for informational purposes only.

Please update your HAN contact information on the Montana Public Health Directory





DATE

March 29, 2024

SUBJECT

Increase in Invasive Serogroup Y Meningococcal Disease in the United States

BACKGROUND

Meningococcal disease, caused by the bacterium *Neisseria meningitidis*, is rare but can result in severe illness with a case-fatality rate of 10–15% even with appropriate antibiotic treatment. Of the six *N. meningitidis* serogroups – A, B, C, W, X, and Y – responsible for most meningococcal disease worldwide, the four serogroups B, C, W, and Y circulate in the United States. Vaccines against serogroups A, C, W, Y (MenACWY) and serogroup B (MenB) are available in the United States. The incidence of *N. meningitidis* serogroup Y, specifically, has increased in the past year in the US.

INFORMATION

There have been no cases of invasive meningococcal disease reported in Montana in the past three years. However, due to increases in this disease in other parts of the country, providers should remain vigilant, especially in populations disproportionately affected by the current increase in the US. Please review the attached HAN for more information.

RECOMMENDATIONS

Clinicians who have a suspect case of meningococcal disease should report it to your [local or tribal health jurisdiction](#) or notify the DPHHS Communicable Disease Epidemiology Section at 406-444-0273. Collect an isolate from a sterile body site such as blood or cerebral spinal fluid. Serogrouping is routinely performed on *N. meningitidis* isolates from sterile body sites at the Montana Public Health Laboratory (MTPHL), and specimens may be stored for epidemiologic purposes. All isolates identified as invasive *Neisseria meningitidis* should be sent to MTPHL for further characterization.

The attached HAN reminds providers that MenACWY vaccines are routinely recommended for adolescents and for people with other risk factors or underlying medical conditions, including HIV. In addition, MTDPHHS reminds providers that MenB vaccination is recommended for certain adolescents and adults with increased risk of disease, such as individuals with functional or anatomic asplenia. All other adolescents may receive the vaccine, preferably at age 16-18.

This is an official
CDC HEALTH ADVISORY

Distributed via the CDC Health Alert Network
March 28, 2024, 1:30 PM ET
CDCHAN-00505

Increase in Invasive Serogroup Y Meningococcal Disease in the United States

Summary

The Centers for Disease Control and Prevention (CDC) is issuing this Health Alert Network (HAN) Health Advisory to alert healthcare providers to an increase in invasive meningococcal disease, mainly attributable to *Neisseria meningitidis* serogroup Y (Figure). In 2023, 422 cases were reported in the United States, the highest annual number of cases reported since 2014. As of March 25, 2024, 143 cases have been reported to CDC for the current calendar year, an increase of 62 cases over the 81 reported as of this date in 2023. A specific meningococcal strain, sequence type (ST) 1466, is responsible for most (101 of 148, 68%) serogroup Y cases with available sequence type data that were reported across the United States in 2023. Cases caused by this strain are disproportionately occurring in people ages 30–60 years (65%), Black or African American people (63%), and people with HIV (15%). In addition, most cases of invasive meningococcal disease caused by ST-1466 in 2023 had a clinical presentation other than meningitis: 64% presented with bacteremia, and at least 4% presented with septic arthritis. Of 94 patients with known outcomes, 17 (18%) died; this case-fatality rate is higher than the historical case-fatality rate of 11% reported for serogroup Y cases in 2017–2021. **Healthcare providers should 1) have a heightened suspicion for meningococcal disease, particularly among populations disproportionately affected by the current increase, 2) be aware that patients may present without symptoms typical of meningitis, and 3) ensure that all people recommended for meningococcal vaccination, including people with HIV, are up to date for meningococcal vaccines.**

Background

[Meningococcal disease](#), caused by the bacterium *Neisseria meningitidis*, is a rare but severe illness with a case-fatality rate of 10–15% even with appropriate antibiotic treatment. Meningococcal disease most often presents as meningitis, with symptoms that may include fever, headache, stiff neck, nausea, vomiting, photophobia, or altered mental status; or as meningococcal bloodstream infection, with symptoms that may include fever and chills, fatigue, vomiting, cold hands and feet, severe aches and pains, rapid breathing, diarrhea, or, in later stages, a dark purple rash. While initial symptoms of meningococcal disease can at first be non-specific, they worsen rapidly, and the disease can become life-threatening within hours. Immediate [antibiotic treatment](#) for meningococcal disease is critical. Survivors may experience long-term effects such as deafness or amputations of the extremities.

Of the six *N. meningitidis* serogroups — A, B, C, W, X, and Y — responsible for most meningococcal disease worldwide, the four serogroups B, C, W, and Y circulate in the United States. Vaccines against serogroups A, C, W, Y (MenACWY) and serogroup B (MenB) are available in the United States. [MenACWY vaccines are routinely recommended](#) for adolescents and for people with other risk factors or underlying medical conditions, including HIV.


Across the United States, 101 ST-1466 cases were reported in 2023. This number is expected to increase with additional laboratory testing data. Cases of invasive meningococcal disease caused by this strain occurred in both males (65%) and females (35%) and disproportionately occurred in people ages 30–60 years (65%), Black or African American people (63%), and people with HIV (15%). In addition, most cases of invasive meningococcal disease caused by ST-1466 had a clinical presentation other than meningitis: 64% presented with bacteremia, and at least 4% presented with septic arthritis. Of 94 patients

with known outcomes, 17 (18%) died; this case-fatality rate is higher than the historical case-fatality rate of 11% reported for serogroup Y cases in 2017–2021.

The serogroup Y ST-1466 strain has contributed to [previously reported increases](#) in meningococcal disease in people with HIV. Based on updated surveillance data, 24 ST-1466 cases have now been reported in people with HIV in 2022–2023; only four were previously vaccinated with MenACWY and none were up to date on recommended doses. To date, no other ST-1466 cases have been identified in people who previously received MenACWY vaccine.

Serogroup Y ST-1466 isolates tested to date have been susceptible to all first-line antibiotics recommended for treatment and prophylaxis. This strain is distinct from [ciprofloxacin-resistant serogroup Y strains that are also circulating in the United States](#) and that are disproportionately affecting Hispanic individuals.

Figure: Trends in meningococcal disease incidence per 100,000 population, by serogroup—United States, 2006–2023. Incidence of *Neisseria meningitidis* serogroup Y is shown in green. Source: National Notifiable Diseases Surveillance System, with additional serogroup data from Active Bacterial Core surveillance (ABCs) and state health departments. 2022 and 2023 data are preliminary.



Recommendations for Public Health Departments

- Ensure healthcare providers are aware of the increasing burden of invasive meningococcal disease, particularly among people ages 30–60 years and Black or African American people.
- Continue submitting all meningococcal isolates to CDC for whole-genome sequencing and antimicrobial susceptibility testing.
- Contact CDC at meningnet@cdc.gov with any questions or concerns about increasing meningococcal disease cases in their jurisdiction or [outbreak](#) investigation or control measures.

Recommendations for Healthcare Providers

- Maintain a heightened suspicion for invasive meningococcal disease and start immediate [antibiotic treatment](#) for persons with suspected meningococcal disease. Blood and cerebrospinal fluid (CSF) cultures are indicated for patients with suspected meningococcal disease [1].
- Recognize that invasive meningococcal disease may affect people of any age or demographic group.
 - Current increases in disease are disproportionately affecting people ages 30–60 years, Black or African American people, and people with HIV.

- Be aware that patients with invasive meningococcal disease may present with bloodstream infection or septic arthritis and **without** symptoms typical of meningitis (e.g., headache, stiff neck).
- Ensure that all people [recommended for meningococcal vaccination](#) are up to date for meningococcal vaccines.
 - All 11–12 year-olds should receive a MenACWY vaccine. Since protection wanes, CDC recommends a booster dose at age 16 years.
 - For people at increased risk due to medical conditions (e.g., with HIV), recommended vaccination includes a 2-dose primary MenACWY series with booster doses every 3–5 years, depending on age.
- Immediately notify [state, tribal, local, or territorial health departments](#) about any suspect or confirmed cases of invasive meningococcal disease.
- Consult with your state or local health department for any questions about meningococcal disease treatment or contact prophylaxis, including any changes based on local meningococcal resistance patterns.

Recommendations for the Public

- Seek medical attention immediately if you or your child develops [symptoms of meningococcal disease](#):
 - Symptoms of meningitis may include fever, headache, stiff neck, nausea, vomiting, photophobia, or altered mental status.
 - Symptoms of meningococcal bloodstream infection may include fever and chills, fatigue, vomiting, cold hands and feet, severe aches and pains, rapid breathing, diarrhea, or, in later stages, a dark purple rash.
 - While symptoms of meningococcal disease can at first be nonspecific, they worsen rapidly, and the disease can become life-threatening within hours.
- Talk to your healthcare provider about [meningococcal vaccines](#) that may be recommended for you and your household or family members, including any recommended booster doses.

For More Information

- Health departments
 - [Meningococcal Disease Surveillance | CDC](#)
 - [Meningococcal Disease | Manual for the Surveillance of Vaccine-Preventable Diseases | CDC](#)
 - [Meningococcal Disease Outbreaks and Public Health Response | CDC](#)
 - [Selection of Antibiotics as Prophylaxis for Close Contacts of Patients with Meningococcal Disease in Areas with Ciprofloxacin Resistance — United States, 2024](#)
- Healthcare providers
 - [Clinical information | Meningococcal Disease | CDC](#)
 - [Meningococcal Vaccination: Information for Healthcare Professionals | CDC](#)
- Everyone
 - [Signs and Symptoms | Meningococcal Disease | CDC](#)
 - [Meningococcal Vaccination | CDC](#)
 - Visit [CDC-INFO](#) or call CDC-INFO at 1-800-232-4636

References

1. American Academy of Pediatrics. Summaries of infectious diseases: meningococcal infections. [Section 3]. In: Kimberlin DW, Barnett ED, Lynfield R, Sawyer MH, eds. Red book: 2021–2024 report of the Committee on Infectious Diseases. Itasca, IL: American Academy of Pediatrics; 2021:519–32.
2. Mbaeyi SA, Bozio CH, Duffy J, et al. Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020. *MMWR Recomm Rep* 2020;69(No. RR-9):1–41. DOI: <http://dx.doi.org/10.15585/mmwr.rr6909a1>

3. Rubis AB, Howie RL, Marasini D, Sharma S, Marjuki H, McNamara LA. Notes from the Field: Increase in Meningococcal Disease Among Persons with HIV — United States, 2022. *MMWR Morb Mortal Wkly Rep* 2023;72:663–664. DOI: <http://dx.doi.org/10.15585/mmwr.mm7224a4>

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages

Health Alert	Conveys the highest level of importance about a public health incident.
Health Advisory	Provides important information about a public health incident.
Health Update	Provides updated information about a public health incident.