Table 11: Tobacco Use, Montana Adults, 2009
(with 95% confidence intervals)

	Current Smokers †				Current Smokeless Tobacco Users ‡			
		95% CI				95% CI		
	Wt.%	LL	UL	UnWt. N	Wt.%	LL	UL	UnWt. N
All Adults	16.8	15.5	18.2	1,268	7.4	6.4	8.5	446
Sex:								
Male	16.4	14.4	18.5	524	13.4	11.6	15.5	385
Female	17.3	15.6	19.1	744	1.5	1.1	2.2	61
Age:								
18 - 24	21.9	15.9	29.3	65	10.2	6.2	16.5	28
25 - 34	23.4	19.2	28.2	143	10.4	7.2	14.7	53
35 - 44	16.0	13.4	18.9	179	9.1	7.1	11.6	96
45 - 54	19.6	17.3	22.1	333	9.0	7.3	11.1	119
55 - 64	14.6	12.8	16.6	302	3.9	2.9	5.2	78
65+	7.8	6.7	9.0	243	3.0	2.3	4.0	70
Education:								
<High School	44.6	37.4	52.1	177	10.1	5.8	17.2	38
High School	21.1	18.5	24.0	505	8.8	7.0	11.0	174
Some College	15.8	13.8	18.1	371	7.5	5.7	9.7	123
College Degree	7.0	5.9	8.4	215	5.3	4.1	6.9	110
Income:								
<$15,000	33.3	28.3	38.8	283	6.4	4.5	9.0	52
$15,000 - $24,999	28.8	24.7	33.3	340	10.5	7.1	15.4	77
$25,000 - $49,999	15.9	13.7	18.4	304	6.9	5.5	8.7	126
$50,000 - $74.999	11.6	9.4	14.3	131	8.0	6.0	10.7	83
$75,000 +	8.8	6.5	11.8	108	6.2	4.6	8.2	72
Race/Ethnicity:								
White, non-Hispanic	15.0	13.7	16.5	967	7.3	6.3	8.4	371
AI/AN*	42.7	36.4	49.3	206	11.7	8.3	16.2	51
Other or Hispanic**	24.8	17.8	33.5	86	6.8	2.8	15.8	22
Disability:								
Disability	22.8	20.1	25.7	438	6.6	5.0	8.6	102
No Disability	15.2	13.8	16.8	818	7.7	6.6	9.1	340
Region:								
1- Eastern MT	15.4	12.9	18.3	172	10.8	8.4	13.8	76
2- N Central MT	20.4	17.6	23.5	272	7.9	6.0	10.2	88
3- S Central MT	18.6	15.6	22.0	192	7.5	5.6	9.9	73
4- Southwest MT	15.8	13.2	18.9	300	7.0	5.3	9.2	106
5- Northwest MT	15.4	13.1	18.1	324	6.8	4.9	9.5	102
MMSA-Billings	18.9	15.0	23.6	109	5.0	3.0	8.3	24
MMSA-Bozeman	10.9	7.1	16.5	55	7.3	4.2	12.5	30
MMSA-Butte	22.7	17.8	28.5	113	4.0	2.6	6.2	26
MMSA-Kalispell	12.6	9.4	16.7	70	4.5	2.1	9.3	16

† A current smoker is defined as someone who has ever smoked 100 cigarettes and who now smokes everyday or some days.
Total Sample Size: 7,596, Weighted Prevalence Estimate: 126,300.

‡ A current user is defined as using chewing tobacco, snuff or snus everyday or some days.
Total Sample Size: 7,615, Weighted Prevalence Estimate: 56,000.

* American Indian or Alaska Native only.

** All other non-White (including multiracial) or Hispanic.