State of Montana *Health Alert Network*

DPHHS HAN

Health INFO SERVICE

Cover Sheet

DATE: June 13, 2016

SUBJECT: Alert to U.S. Healthcare Facilities: First *mcr-1* Gene in *E. coli* Bacteria found in a Human in the United States

For LOCAL HEALTH DEPARTMENT Reference Only

DPHHS Subject Matter Resource for more information regarding this HAN, contact:

DPHHS CDCP
Epidemiology Section
1-406-444-0273

INSTRUCTIONS:

DISTRIBUTE AT YOUR DISCRETION. Share this information with relevant SMEs or contacts (internal and external) as you see fit.

DPHHS Health Alert Hotline:
1-800-701-5769

DPHHS HAN Website:
www.han.mt.gov

REMOVE THIS COVER SHEET BEFORE REDISTRIBUTING AND REPLACE IT WITH YOUR OWN

Please ensure that DPHHS is included on your HAN distribution list.
hhshan@mt.gov

Categories of Health Alert Messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.
Health Advisory: provides important information for a specific incident or situation; may not require immediate action.
Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.
Information Service: passes along low level priority messages that do not fit other HAN categories and are for informational purposes only.

Please update your HAN contact information on the Montana Public Health Directory

State of Montana ***Health Alert Network***

DPHHS HAN

Information Sheet

Date: June 13, 2016

Subject: Alert to U.S. Healthcare Facilities: First *mcr-1* Gene in *E. coli* Bacteria found in a Human in the United States

Background: N/A

Information: N/A

Recommendations: Read the attached CDC HAN message.

This is an official

CDC HEALTH ADVISORY

Distributed via the CDC Health Alert Network
June 13, 2016, 13:35 EDT (1:35 PM EDT)
CDCHAN-00390

Alert to U.S. Healthcare Facilities: First *mcr-1* Gene in *E. coli* Bacteria found in a Human in the United States

Summary
The Centers for Disease Control and Prevention (CDC) is collaborating in a coordinated public health response to the Department of Defense (DoD) announcement on May 26 of the first *mcr-1* gene found in bacteria from a human in the United States (http://aac.asm.org/content/early/2016/05/25/AAC.01103-16.full.pdf+html). *Escherichia coli* (*E. coli*) bacteria carrying the *mcr-1* gene were found in a urine sample from a person in Pennsylvania with no recent travel outside of the United States who presented to a clinic with a urinary tract infection. The *mcr-1* gene makes bacteria resistant to the antibiotic colistin, which is used as a last-resort drug to treat patients with infections caused by multidrug-resistant bacteria, including carbapenem-resistant Enterobacteriaceae (CRE). The *mcr-1* gene exists on a plasmid, a small piece of DNA that is capable of moving from one bacterium to another, potentially spreading antibiotic resistance to other bacterial species. CDC is issuing this HAN notice as a reminder to U.S. healthcare facilities about recommendations to prevent antibiotic resistant infections and alert them to additional recommendations for detecting and reporting bacteria with the *mcr-1* gene.

Background
In November 2015, a report from China first described plasmid-mediated colistin-resistance caused by the *mcr-1* gene. Following that report, retrospective investigations of historical isolates from outside the United States have identified the rare occurrence of *mcr-1* in Enterobacteriaceae from the 1980s. Bacteria with this resistance mechanism have now been identified from humans, food, environmental samples, and food animals in at least 20 countries around the world. Most reports to date have identified the *mcr-1* gene in *E. coli*, but it has also been reported from *Salmonella* species, *Shigella sonnei*, and *Klebsiella pneumoniae*.

Three *mcr-1* producing *E. coli* have been identified in the United States as of June 7, 2016: one in a clinical specimen from a person in Pennsylvania and two from intestinal samples from pigs. The *E. coli* isolate from the patient was also resistant to antibiotics in at least five additional antibiotic classes, including cephalosporins, fluoroquinolones, sulfonamides, aminoglycosides, and tetracyclines. The sample from one pig also was resistant to other antibiotics, including ampicillin, streptomycin, sulfisoxazole, and tetracycline. No additional resistance was found in the isolate from the second pig. The presence of the *mcr-1* gene on a plasmid means that colistin resistance can be shared with other more resistant bacteria such as CRE, raising the possibility that untreatable bacteria could develop. A rapid public health response is underway to identify and contain any potential spread from the patient. CDC laboratories have developed protocols for testing microorganisms for the *mcr-1* gene and are performing screening tests to see if people in contact with the patient with *mcr-1* might be colonized with this organism. CDC is increasing its surveillance of human samples from U.S. healthcare settings. CDC's National Antimicrobial Resistance Monitoring System, in collaboration with the Food and Drug Administration (FDA) and the U.S. Department of Agriculture (USDA), will continue to look for *mcr-1* mediated colistin resistance in enteric bacteria from humans, retail meat, and food animals.

Recommendations
Given the discovery of *mcr-1* in a person in Pennsylvania, CDC reiterates the importance of measures to

prevent transmission of antibiotic resistant bacteria, including those resistant to colistin or carrying the *mcr-1* gene. CDC recommends the following:

- **Infection Prevention:** Healthcare providers should follow Standard and Contact Precautions (http://www.cdc.gov/hicpac/2007IP/2007ip_part3.html) for any patients colonized or infected with antibiotic resistant bacteria, including patients who are found to have *mcr-1* mediated resistant organisms. Healthcare facilities should follow manufacturers' instructions for device cleaning and reprocessing.
- **Laboratory Testing:** If laboratories are testing to determine whether colistin can be used clinically, Enterobacteriaceae isolates with a minimum inhibitory concentration (MIC) to colistin of 4 µg/ml or higher should be tested for confirmation and the presence of *mcr-1*. Thus far, all microorganisms that have contained the *mcr-1* gene can safely be tested in a biosafety level-2 (BSL-2) laboratory. Isolates should be sent to CDC for confirmatory testing via the state or local public health department, per the CDC test directory (http://www.cdc.gov/laboratory/specimen-submission/detail.html?CDCTestCode=CDC-10223), if local testing is not available. The results and test method that were used for initial colistin testing should be included with any isolates submitted for confirmatory testing. CDC laboratories are in the process of validating a rapid polymerase chain reaction (PCR) test to detect *mcr-1* in bacteria with elevated colistin MICs. It is not necessary to test Enterobacteriaceae with intrinsic colisitin resistance (e.g., *Proteus*, *Providencia*, *Morganella*, and *Serratia* species). Additionally, since *Enterobacter* species often have MICs of >=2 mcg/ml to colisitin, they should be sent for *mcr-1* testing only if other risk factors exist, such as a recent history of travel outside the United States to a country where *mcr-1* has been found to be more common (http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=21403).
- **Validation of Laboratory Testing:** CDC is making test-bacteria with elevated colistin MICs, available to laboratories, researchers, and others through the FDA-CDC Antimicrobial Resistance Bacteria Isolate Bank (http://www.cdc.gov/drugresistance/resistance-bank/) for use in validation of colistin-resistance testing in U.S. clinical laboratories.
- **Environmental Cleaning**: Healthcare facilities should ensure rooms where patients with antibiotic-resistant infections have been placed receive thorough daily and terminal cleaning.
- **Reporting to Public Health:** Healthcare facilities and laboratories should adhere to local reporting requirements for all antibiotic resistant infections. If Enterobacteriaceae with *mcr-1* are identified from patients, healthcare facilities and laboratories should notify local or state public health authorities as quickly as possible, and inform clinicians caring for the patient and responsible infection prevention staff.
- **Preparing food safely:** Cook all meat, poultry, and fish to its proper internal temperature to kill bacteria **(**http://www.foodsafety.gov/keep/charts/mintemp.html**),** viruses, and other foodborne pathogens, regardless of antibiotic resistance.

For More Information

1. CDC Facility Guidance for Control of CRE: November 2015 Update: http://www.cdc.gov/hai/pdfs/cre/CRE-guidance-508.pdf
2. Responding to new Forms of Antibiotic Resistance: http://www.cdc.gov/hai/pdfs/toolkits/Responding-to-New-Forms-of-Antibiotic-Resistance.pdf
3. Biosafety in Microbiological and Biomedical Laboratories, 5th Edition: http://www.cdc.gov/biosafety/publications/bmbl5/BMBL.pdf
4. CDC Vital Signs Report: Making Health Care Safer, Protect Patients from Antibiotic Resistance: http://www.cdc.gov/vitalsigns/protect-patients/index.html

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages:
Health Alert Requires immediate action or attention; highest level of importance
Health Advisory May not require immediate action; provides important information for a specific incident or situation
Health Update Unlikely to require immediate action; provides updated information regarding an incident or situation
HAN Info Service Does not require immediate action; provides general public health information

##This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations##