



Montana

Risk Factors and Health Indicators Report

Summary Table of Risk Factors and Health Indicators in Montana

	Indicator	State	National
Death Rates	Motor Vehicle Death Rate (2012) Fatality rate per 100,000 population	20.4	10.69
	Infant Mortality Rate (2011) Number of infant deaths (before age 1) per 1,000 live births	5.88	6.07
	Homicide Death Rate (2011) Age-adjusted deaths due to homicide/assault (per 100,000 population)	2.8	5.3
	Stroke Death Rate (2011) Age adjusted rate of stroke deaths per 100,000 population	37.9	37.9
	Suicide Death Rate (2011) Age-adjusted deaths due to suicide/intentional self-harm (per 100,000 population)	22.6	12.5
	Drug Poisoning Death Rate (2011) Rate of drug poisoning deaths per 100,000 population	14.4	13.2
	Heart Disease Death Rate (2011) Age adjusted rate of heart disease deaths per 100,000 population	156.9	173.7
	Cancer Death Rate (2011) Age-adjusted rate of cancer deaths per 100,000 population	164.8	169
Health Burden	CLABSI-Standardized Infection Ratio (2012) Central Line-Associated Bloodstream Infections -Standardized Infection Ratios	0.488	0.563
	Hepatitis B Cases (2011) Number of new cases of acute hepatitis B (per 100,000 population)	0	2890
	Hepatitis C Cases (2011) Number of new cases of acute hepatitis C (per 100,000 population)	9	1229
	Diagnosed Diabetes (2012) Percent of adults (age 18+) ever told by health professional that they have diabetes	7.2	9.7
	Teen Birth Rate (2012) Birth Rate for teens ages 15-19 per 1,000 female population.	35	34.2
	Youth Obesity (2013) Percent of high school students who are obese (BMI > the 95th percentile for age/sex)	9.4%	13.7%
	HIV Diagnosis Rate (2011) Rate of persons diagnosed with HIV infection per 100,000 population	2.6	19
	Adult Obesity (2012) Percent of Adults (age 18+) who are obese (BMI > 30)	24.3	28.1
	Diagnosed High Cholesterol (2011) Percent of adults who have had their blood cholesterol checked and have been told it was high	34.6%	38.4%
	Diagnosed Hypertension (2011) Percent of Adults (age 18+) ever told by health professional you have High blood pressure	30.2%	30.9%
	Medicated Hypertension (2011) Percent of Adults (age 18+) currently taking medicine for high blood pressure (hypertension)	73.1%	77.7%



Centers for Disease Control and Prevention Sortable Stats - Risk Factors and Health Indicators

	Indicator	State	National
Risk Factors	Observed Seat Belt Use (2012) Percent of observed seat belt use based on probability samples in all 50 states	76.9	86
	Youth Nutrition (2013) Percent of high school students that ate vegetables less than one time per day	33.2%	38.5%
	Youth Binge Drinking (2013) Percent of high school students who reported having five or more drinks of alcohol in a row within a couple of hours on at least 1 day in last 30 days	23.5%	20.8%
	Youth Marijuana Use (2013) Percent of high school students who used marijuana on at least 1 or more days in the last 30 days	21	23.4
	Adult Binge Drinking (2012) Percent of Adults (age 18+) who are binge drinkers (males having five or more drinks on one occasion, females having four or more drinks on one occasion)	21.8	16.9
	Adult Smoking (2012) Percent of Adults (age 18+) who currently smoke cigarette	19.7	19.6
	Youth Physical Activity (2013) Percent of high school student that are physically active at least 60 minutes per day on five or more days	54.8%	47.3%
	Youth Seat Belt Use (2013) Percent of high school students who wore a seat belt sometimes, most of the time, or always	89.9%	92.4%
	Youth Smoking (2013) Percent of high school students who smoked cigarettes on at least 1 or more days in last 30 days	13.2%	15.7%
	Adult Physical Activity (2011) Percent of adults that participated in 150 minutes or more of Aerobic Physical Activity per week	55.3%	51.6%
	Adult Nutrition (2009) Percent of adults that consume fruits and vegetables five or more times per day	25.7%	23.5%
	Preventive Services	Child Vaccination Coverage (2013) Percent of children aged 19 to 35 months receiving the recommended doses of DTaP, polio, MMR, Hib, hepatitis B, varicella and PCV.	65.4
Colorectal Cancer Screening (2012) Percent of persons age 50+ who have ever had a sigmoidoscopy or colonoscopy		0.665	0.668
Influenza Vaccination Coverage (2012) Percent of persons aged 6 months and older who received a seasonal influenza vaccination within the past year		41.7%	45%

Suggested Citation: Centers for Disease Control and Prevention. Sortable Risk Factors and Health Indicators Website, last updated (April, 2013). Site accessed at <http://www.cdc.gov/sortablestats/>.

This report was generated from CDC Sortable Stats web application, an interactive data set comprised of behavioral risk factors and health indicators compiled from various published CDC and federal sources. To access the application, please visit <http://www.cdc.gov/sortablestats/>. Refer to the About the Data section of the application for the detailed information about indicators, data sources and for additional resources.

(http://www.cdc.gov/sortablestats/Report_Docs/PDFDocs/Sortable_Stats_Data_Sources.pdf)