

# Interview Periods Resource

Disease Code	Disease Type	Interview Period
200/300	Chlamydia/Gonorrhea	<p>Symptomatic—60 days prior to onset of symptoms through the date of treatment</p> <p>Asymptomatic cases—60 days prior to initial diagnostic test</p>
490	Pelvic Inflammatory Disease	60 days prior to onset of symptoms through the date of treatment
710	Primary Syphilis	90 days prior to the onset of symptoms up to the date of adequate treatment
720	Secondary Syphilis	6½ months prior to the onset of symptoms up to the date of adequate treatment
730	Early Latent Syphilis	12 months prior to the date of adequate treatment
900/950	HIV/AIDS	<p>Use local protocols, if none exist:</p> <p>1 year prior to the date that the positive specimen was collected through the date of the interview (extended interview period may be warranted by individual circumstances)</p> <p>6 months prior to a negative test date [through the date of the interview] (With more recent testing technology, it may be possible to shorten the 3 months prior to the last negative test date.)</p> <p>10-year interview period for current or any previous spouses</p>



## Interview Periods Resource

Disease	Also Consider
Gonorrhea/Chlamydia	If no partner is found within 60 days, go back to the last known partner for Chlamydia
Pelvic Inflammatory Disease	If no partner is found within 60 days, go back to the last known partner
Syphilis	If a negative test is confirmed, the interview period begins 97 days prior to the negative test.  A history of primary symptoms on a secondary patient or secondary symptoms on an early latent patient: use symptoms to determine the start of the interview period.
HIV	6 months prior to the last negative test (may be shortened to 3 months with some of the newer testing technology)