

DPHHS HAN ADVISORY



Cover Sheet

DATE: January 28, 2013

SUBJECT: Guidance for Shortages of Isoniazid (INH)
and Treatment for Tuberculosis

INSTRUCTIONS:

DISTRIBUTE to your local HAN contacts. This HAN is intended for general sharing of information. **Remove this cover sheet before redistributing and replace it with your own.**

For LOCAL HEALTH DEPARTMENT reference only

DPHHS Subject Matter Resource for more information regarding this HAN, contact:

**Communicable Disease &
Prevention Bureau
TB/Epidemiology Section
406-444-0273**

**DPHHS Health Alert Hotline:
1-800-701-5769**

**DPHHS HAN Website:
www.han.mt.gov**

Remove this cover sheet before redistributing and replace it with your own.

Please ensure that DPHHS is included on your HAN distribution list.

Categories of Health Alert Messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.

Please call DPHHS to update contact information at 444-0919

Information Sheet

Date: January 28, 2013

Subject: Recommendations for Drug Allocation, Tuberculosis (TB) Prevention, and Patient Care during Isoniazid (INH) Shortages

Information: Please see the attached CDC Health Alert Network (HAN) Advisory regarding how to adjust practices in response to emerging shortages of INH.

Key Points & Recommendations:

- The DPHHS Montana Tuberculosis Program is taking steps to ensure the supply of Isoniazid (INH) currently available at the state contract pharmacy is preserved for the highest priority patients
- The state program is generally used as a “payer of last resort” for treating active cases of TB. If the supply of INH at private or governmental hospitals/clinics is depleted, the state supply might need to be used for patients who normally would be eligible to obtain INH through other means, such as private health insurance, the Indian Health Service, Tribal Health, the Veterans Administration, etc.
- Please check your current private supply and continue to use your existing suppliers when possible

Clinical and Public Health Recommendations:

As of today, the following actions are being implemented by the Montana TB Program:

- 1) Most of the current supply of INH at the DPHHS contract pharmacy will be reserved for existing or new active cases of tuberculosis.
- 2) All INH orders normally supplied through the DPHHS contract pharmacy for either active cases or patients with latent TB infection (LTBI) must be pre-approved by the Montana Tuberculosis Program.
- 3) **No new INH orders** through the DPHHS contract pharmacy will be filled for patients with LTBI unless the patient is high-risk of developing disease, i.e. patients diagnosed with LTBI during a contact investigation, children <5 years old with LTBI, and HIV-infected patients with LTBI. Requests for other at-risk patients will be determined on a case-by-case basis. For new patients being considered for treatment of LTBI, rifampin administered for 4 months can be used for most patients.
- 4) For LTBI patients already being treated with a 6- or 9-month course of INH, please contact the Montana TB Program for assistance in either changing to the 4-month rifampin regimen or for assistance in securing a supply of INH to complete the prescribed course.

For more information, contact Denise Ingman at dingman@mt.gov or 406-444-0273.

This is an official **CDC HEALTH ADVISORY**

Distributed via the CDC Health Alert Network
January 28, 2013, 09:00 ET
CDCHAN-00340

Recommendations for Drug Allocation, Tuberculosis Prevention, and Patient Care During Isoniazid Shortages

Summary: Shortages of isoniazid (INH), a cornerstone drug for treating tuberculosis disease (TB) and latent *Mycobacterium tuberculosis infection* (LTBI), are continuing. This notice gives an update on the shortages and expands general guidance to public health officials and clinicians about how to adjust practices in response to the shortages from that outlined in CDC's December 21, 2012, *Morbidity and Mortality Weekly Report* (MMWR) (<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6150a4.htm>). It also outlines national plans for restoring INH supplies and lists published guidance that could assist in making treatment decisions when INH is unavailable.

Background

U.S. INH production has been interrupted, and stocks are dwindling. Consequent to the critical shortage of 300 mg tablets that was recently reported (1), the stocks of 100 mg tablets in some localities are reportedly being depleted. An initial forecast date for restoring INH production has been extended from late January 2013 to as late as March 2013 (1,2); however, the forecasts are unstable and vary by company. One of three pharmaceutical companies supplying INH, VersaPharm Incorporated, Marietta, Georgia, has notified public health officials that it is cancelling all INH backorders and not accepting new orders.

Clinical and Public Health Recommendations

While INH is in shortage, clinicians should coordinate TB and LTBI treatment plans with their jurisdictional public health officials, who should set priorities for allocating INH stock locally. These priorities should be set according to the immediate medical needs of individual patients and a general TB control strategy:

- 1) treating patients who have TB disease,
- 2) treating LTBI patients who are diagnosed during contact tracing of contagious TB (i.e., contact investigations), and
- 3) treating LTBI patients who face the greatest likelihood of TB disease (e.g., HIV-infected patients) or the hazard of severe illness (e.g., children <5 years old).

Treatment for LTBI can be postponed when the likelihood or hazard of TB disease is low, but systems should be developed for recalling patients when INH becomes available.

The World Health Organization (WHO) recommends that "drugs for the full course of treatment are reserved for the patient at the outset of treatment" (3). However, the abrupt onset of the INH shortage means that treatment for some TB patients could be disrupted in mid-course. Decisions for continuing treatment in these situations should be made on a case-by-case basis in consultation with a TB expert, as arranged by jurisdictional public health officials.

Fixed dose combination formulations of INH and rifampin or INH, rifampin, and pyrazinamide can be used for treating TB disease if the dosages of all the drugs are correct for the patient (4). However, at least one fixed dose combination is already on backorder, and stocks are believed to be depleted. A full treatment course of drugs should be secured before starting treatment with one of these combinations.

Regimens without INH for treating TB disease are less well studied than those with INH. CDC, the American Thoracic Society, and the Infectious Diseases Society of America have recommended a regimen of rifampin, pyrazinamide, and ethambutol for 6 months, with consideration of a fluoroquinolone, such as moxifloxacin or levofloxacin, for more extensive disease, when the infecting *M. tuberculosis* isolate is INH-resistant or the patient is intolerant of INH (4). Directly observed therapy should be used for reducing the potential for acquired rifampin resistance.

An alternative treatment regimen for LTBI, when the infecting *M. tuberculosis* is believed to be INH resistant, or when the patient cannot tolerate INH, is rifampin administered for 4 months, either self-administered or directly observed (5). This regimen has been adopted for routine use by some U.S. TB program directors. A combination regimen of INH and rifampin for 3 or 4 months is used overseas and in some U.S. jurisdictions, although it is not currently recommended in U.S. national guidelines (5,6). This regimen could be prescribed via a fixed dose combination of INH and rifampin if supplies for complete treatment courses are in stock. The combination regimen of rifampin and pyrazinamide for 2 months is *not recommended because of excess toxicity*. Any of the alternative regimens for TB disease or LTBI could be more expensive than the standard regimens.

CDC is working with the Food and Drug Administration (FDA), the National Tuberculosis Controllers Association, and the two pharmaceutical companies currently planning to supply INH (Teva Pharmaceuticals USA, Sellersville, Pennsylvania; and Sandoz Inc., Princeton, New Jersey) to determine current INH stocks, jurisdictions with urgent needs, and options for restoring supplies as soon as can be safely done. CDC will monitor notifications from public health officials of INH shortages and stocks, and will provide updates as they are available on the website for the Division of Tuberculosis Elimination at <http://www.cdc.gov/tb/>.

References for More Information:

1. CDC. Notes from the field: national shortage of isoniazid 300 mg tablets. *MMWR*. 2012;61:1029. Available at http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6150a4.htm?s_cid=mm6150a4_w
2. FDA. Drug safety and availability. Available at <http://www.fda.gov/drugs/drugsafety/drugshortages/default.htm>.
3. World Health Organization. Treatment of tuberculosis: guidelines, fourth edition. WHO Press, 2010. Available at http://whqlibdoc.who.int/publications/2010/9789241547833_eng.pdf.
4. CDC. Treatment of Tuberculosis, American Thoracic Society, CDC, and Infectious Diseases Society of America. *MMWR* 2003;52(No. RR-11). Available at <http://www.cdc.gov/mmwr/PDF/rr/rr5211.pdf>.
5. CDC. Targeted tuberculin testing and treatment of latent tuberculosis infection. *MMWR* 2000;49(No. RR-6), available at <http://www.cdc.gov/mmwr/PDF/rr/rr4906.pdf>.
6. National Institute for Health and Clinical Excellence. Tuberculosis: clinical diagnosis and management of tuberculosis, and measures for its prevention and control. NICE clinical guideline 117. Royal College of Physicians of London, 2006 and 2011. Available at <http://www.nice.org.uk/nicemedia/live/13422/53638/53638.pdf>.

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages:

Health Alert Requires immediate action or attention; highest level of importance

Health Advisory May not require immediate action; provides important information for a specific incident or situation

Health Update Unlikely to require immediate action; provides updated information regarding an incident or situation

HAN Info Service Does not require immediate action; provides general public health information

##This message was distributed to state and local health officers, public information officers, epidemiologists, HAN coordinators, and clinician organizations##