

Montana Health Alert Network

DPHHS HAN INFO SERVICE

Cover Sheet

DATE

June 11, 2021

SUBJECT

Increased Interseasonal Respiratory Syncytial Virus (RSV) Activity in Parts of the Southern United States

INSTRUCTIONS

DISTRIBUTE AT YOUR DISCRETION. Share this information with relevant SMEs or contacts (internal and external) as you see fit.



For LOCAL HEALTH
DEPARTMENT reference only
DPHHS Subject Matter Resource for
more information regarding this HAN,
contact:

**Epidemiology Section
1-406-444-0273**

*For technical issues related to the HAN
message contact the Emergency
Preparedness Section
at 1-406-444-0919*

DPHHS Health Alert Hotline:
1-800-701-5769

DPHHS HAN Website:
www.han.mt.gov

REMOVE THIS COVER SHEET BEFORE REDISTRIBUTING AND REPLACE IT WITH YOUR OWN

Please ensure that DPHHS is included on your HAN distribution list.
hhshan@mt.gov

Categories of Health Alert Messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.

Information Service: passes along low level priority messages that do not fit other HAN categories and are for informational purposes only.

Please update your HAN contact information on the Montana Public Health Directory

DPHHS HAN

Information Sheet



DATE

June 11, 2021

SUBJECT

Increased Interseasonal Respiratory Syncytial Virus (RSV) Activity in Parts of the Southern United States

BACKGROUND

RSV is an RNA virus of the genus *Orthopneumovirus*, family *Pneumoviridae*, primarily spread via respiratory droplets when a person coughs or sneezes, and through direct contact with a contaminated surface. RSV is the most common cause of bronchiolitis and pneumonia in children under one year of age in the United States. Infants, young children, and older adults with chronic medical conditions are at risk of severe disease from RSV infection. Each year in the United States, RSV leads to on average approximately 58,000 hospitalizations¹ with 100-500 deaths among children younger than 5 years old² and 177,000 hospitalizations with 14,000 deaths among adults aged 65 years or older.

In Montana, the RSV season typically extends from late fall through early spring. This year, over 9,000 RSV tests were completed by local laboratories but only one positive was identified during the during this time, which is much less than expected. However, over the past six weeks very localized RSV activity has been reported in the Lewis and Clark County region.

INFORMATION

The Centers for Disease Control and Prevention (CDC) is issuing this health advisory to notify clinicians and caregivers about increased interseasonal respiratory syncytial virus (RSV) activity across parts of the Southern United States. Due to this increased activity, CDC encourages broader testing for RSV among patients presenting with acute respiratory illness who test negative for SARS-CoV-2, the virus that causes COVID-19. RSV can be associated with severe disease in young children and older adults. This health advisory also serves as a reminder to healthcare personnel, childcare providers, and staff of long-term care facilities to avoid reporting to work while acutely ill – even if they test negative for SARS-CoV-2

RECOMMENDATIONS

See the attached CDC HAN.

This is an official
CDC HEALTH ADVISORY

Distributed via the CDC Health Alert Network
June 10, 2021, 1:30 ET PM
CDCHAN-00443

Increased Interseasonal Respiratory Syncytial Virus (RSV) Activity in Parts of the Southern United States

Summary

The Centers for Disease Control and Prevention (CDC) is issuing this health advisory to notify clinicians and caregivers about increased interseasonal respiratory syncytial virus (RSV) activity across parts of the Southern United States. Due to this increased activity, CDC encourages broader testing for RSV among patients presenting with acute respiratory illness who test negative for SARS-CoV-2, the virus that causes COVID-19. RSV can be associated with severe disease in young children and older adults. This health advisory also serves as a reminder to healthcare personnel, childcare providers, and staff of long-term care facilities to avoid reporting to work while acutely ill – even if they test negative for SARS-CoV-2.

Background

RSV is an RNA virus of the genus *Orthopneumovirus*, family *Pneumoviridae*, primarily spread via respiratory droplets when a person coughs or sneezes, and through direct contact with a contaminated surface. RSV is the most common cause of bronchiolitis and pneumonia in children under one year of age in the United States. Infants, young children, and older adults with chronic medical conditions are at risk of severe disease from RSV infection. Each year in the United States, RSV leads to on average approximately 58,000 hospitalizations¹ with 100-500 deaths among children younger than 5 years old² and 177,000 hospitalizations with 14,000 deaths among adults aged 65 years or older.³

In the United States, RSV infections occur primarily during the fall and winter cold and flu season. In April 2020, RSV activity decreased rapidly, likely due to the adoption of public health measures to reduce the spread of COVID-19.⁴ Compared with previous years, RSV activity remained relatively low from May 2020 to March 2021. However, since late March, CDC has observed an increase in RSV detections reported to the [National Respiratory and Enteric Virus Surveillance System](#) (NREVSS), a nationwide passive, laboratory-based surveillance network. CDC noted increases in laboratory detections and in the percentages of positive detections for both antigen and PCR testing in parts of HHS Region 4 (Alabama, Florida, Georgia, Kentucky, Mississippi, North Carolina, South Carolina, and Tennessee) and Region 6 (Arkansas, Louisiana, New Mexico, Oklahoma, and Texas). Due to limited testing outside of the typical RSV season, data are limited in some jurisdictions and may be incomplete for the most recent weeks. Since this elevated interseasonal activity is a deviation in the typical circulation patterns for RSV, at this time it is not possible to anticipate the likely spread, peak, or duration of activity with any certainty. Health officials also identified increased interseasonal RSV circulation in parts of Australia during late 2020 and in South Africa in early 2021. Still, RSV did not reach seasonal peak levels in most regions or result in widespread circulation.⁵⁻⁷

Due to reduced circulation of RSV during the winter months of 2020–2021, older infants and toddlers might now be at increased risk of severe RSV-associated illness since they have likely not had typical levels of exposure to RSV during the past 15 months. In infants younger than six months, RSV infection may result in symptoms of irritability, poor feeding, lethargy, and/or apnea with or without fever. In older infants and young children, rhinorrhea and decreased appetite may appear one to three days before cough, often followed by sneezing, fever, and sometimes wheezing. Symptoms in adults are typically consistent with upper respiratory tract infections, including rhinorrhea, pharyngitis, cough, headache, fatigue, and fever. There is no specific treatment for RSV infection other than [symptom management](#).

Recommendations

1. Clinicians and caregivers should be aware of the typical clinical presentation of RSV for different age groups.
2. Clinicians should consider testing patients with a negative SARS-CoV-2 test and acute respiratory illness or the age-specific symptoms presented above for non-SARS-CoV-2 respiratory pathogens, such as RSV. Real-time reverse transcription-polymerase chain reaction (rRT-PCR) is the preferred method for testing for respiratory viruses.
3. Clinicians should report laboratory-confirmed RSV cases and suspected clusters of severe respiratory illness to local and state health departments according to their routine reporting requirements.
4. Healthcare personnel, childcare providers, and staff of long-term care facilities should avoid reporting to work while acutely ill – even if they test negative for SARS-CoV-2.
5. Clinicians can review weekly updates to the [NREVSS website](#) and refer to surveillance data collected by local hospitals and health departments for information on RSV circulation trends in their area.

For More Information

- [CDC – RSV Information for Healthcare Providers](#)
- [CDC – RSV National Trends - NREVSS](#)
- [CDC – RSV Trends and Surveillance](#)
- [CDC – RSV Symptoms and Care](#)

References

¹[Rha B, Curns AT, Lively JY, Campbell AP, Englund JA, Boom JA, Azimi PH, Weinberg GA, Staat MA, Selvarangan R, Halasa NB, McNeal MM, Klein EJ, Harrison CJ, Williams JV, Szilagyi PG, Singer MN, Sahni LC, Figueroa-Downing D, McDaniel D, Prill MM, Whitaker BL, Stewart LS, Schuster JE, Pahud BA, Weddle G, Avadhanula V, Munoz FM, Piedra PA, Payne DC, Langley G, Gerber SI. Respiratory Syncytial Virus-Associated Hospitalizations Among Young Children: 2015-2016. *Pediatrics*. 2020 Jul;146\(1\):e20193611. doi: 10.1542/peds.2019-3611. Epub 2020 Jun 16. PMID: 32546583.](#)

²[Thompson WW, Shay DK, Weintraub E, Brammer L, Cox N, Anderson LJ, Fukuda K. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA*. 2003 Jan 8;289\(2\):179-86. doi: 10.1001/jama.289.2.179. PMID: 12517228.](#)

³[Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. Respiratory syncytial virus infection in elderly and high-risk adults. *N Engl J Med*. 2005 Apr 28;352\(17\):1749-59. doi: 10.1056/NEJMoa043951. PMID: 15858184.](#)

⁴[Haddadin Z, Schuster JE, Spieker AJ, Rahman H, Blozinski A, Stewart L, Campbell AP, Lively JY, Michaels MG, Williams JV, Boom JA, Sahni LC, Staat M, McNeal M, Selvarangan R, Harrison CJ, Weinberg GA, Szilagyi PG, Englund JA, Klein EJ, Curns AT, Rha B, Langley GE, Hall AJ, Patel MM, Halasa NB. Acute Respiratory Illnesses in Children in the SARS-CoV-2 Pandemic: Prospective Multicenter Study. *Pediatrics*. 2021 May 13:e2021051462. doi: 10.1542/peds.2021-051462. Epub ahead of print. PMID: 33986150.](#)

⁵[Foley DA, Yeoh DK, Minney-Smith CA, Martin AC, Mace AO, Sikazwe CT, Le H, Levy A, Moore HC, Blyth CC. The Interseasonal Resurgence of Respiratory Syncytial Virus in Australian Children Following the Reduction of Coronavirus Disease 2019-Related Public Health Measures. *Clin Infect Dis*. 2021 Feb 17:ciaa1906. doi: 10.1093/cid/ciaa1906. Epub ahead of print. PMID: 33594407; PMCID: PMC7929151.](#)

⁶[Virus Watch](#), Week Ending 30 May 2021. Government of Western Australia, Department of Health. Accessed 6/7/2021.

⁷[Weekly Respiratory Pathogens Surveillance Report](#), South Africa, Week 21, 2021. National Institute for Communicable Diseases, Division of the National Health Laboratory Service. Accessed 6/7/21.

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages:

Health Alert Requires immediate action or attention; highest level of importance

Health Advisory May not require immediate action; provides important information for a specific incident or situation

Health Update Unlikely to require immediate action; provides updated information regarding an incident or situation

HAN Info Service Does not require immediate action; provides general public health information

##This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations##