

Montana Health Alert Network

DPHHS HAN

ADVISORY

Cover Sheet

DATE

March 19, 2024

SUBJECT

Increase in Global and Domestic Measles Cases and Outbreaks: Ensure Children in the United State and Those Traveling Internationally 6 Months and Older are Current on MMR Vaccination

INSTRUCTIONS

DISTRIBUTE to your local HAN contacts. This HAN is intended for general sharing of information.

- Time for Forwarding: **As Soon As Possible**
- Please forward to DPHHS at hhshan@mt.gov
- **Remove this cover sheet before redistributing and replace it with your own**



For LOCAL HEALTH DEPARTMENT reference only
DPHHS Subject Matter Resource for more information regarding this HAN,

contact:

DPHHS CDCP

Epidemiology Section
1-406-444-0273

Immunization Section
1-406-444-5580

For technical issues related to the HAN message contact the Emergency Preparedness Section at 1-406-444-0919

DPHHS HAN Website:
<https://dphhs.mt.gov/publichealth/han>

REMOVE THIS COVER SHEET BEFORE REDISTRIBUTING AND REPLACE IT WITH YOUR OWN

Please ensure that DPHHS is included on your HAN distribution list.
hhshan@mt.gov

Categories of Health Alert Messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.

Information Service: passes along low level priority messages that do not fit other HAN categories and are for informational purposes only.

Please update your HAN contact information on the Montana Public Health Directory



DATE

March 19, 2024

SUBJECT

Increase in Global and Domestic Measles Cases and Outbreaks: Ensure Children in the United States and Those Traveling Internationally 6 Months and Older are Current on MMR Vaccination

BACKGROUND

Measles is a highly contagious, vaccine-preventable illness that remains a public health concern. Cases of measles are on the rise both globally and domestically. Since December 2023, the U.S. has seen a sharp increase in cases related to unvaccinated travelers exposed while abroad who becoming symptomatic after returning home. This can lead to an outbreak if unvaccinated persons are exposed. As of March 14, 2024, there have been 58 confirmed cases of measles in 17 states since the start of 2024. As of March 19, 2024, there have been no confirmed cases of measles in Montana.

INFORMATION

Measles is a viral illness that spreads to 9 out of 10 exposed unvaccinated close contacts. Cases of measles have been reported in unvaccinated travelers returning to the US, particularly in unvaccinated children. The last case of measles reported in Montana was in 1990.

RECOMMENDATIONS

Cases of measles are immediately reportable in the US and Montana. If a patient is suspected to have measles, please immediately notify your [local or tribal health department](#). If you are unable to contact the local public health department, please call Montana DPHHS at 406-444-0273. This number is available 24/7. DPHHS can help coordinate testing for suspect measles cases at the state public health laboratory, provide infection control recommendations, and assist local public health with case investigations and contact tracing. Please see the included attachment for additional measles testing guidance at the Montana Public Health Laboratory.

Healthcare providers should review MMR routine and international travel vaccination recommendations in the attached CDC HAN. International travel vaccine recommendations may differ from routine vaccination schedules. MMR vaccine can be found at local and tribal health departments, healthcare provider offices and many pharmacies.

This is an official
CDC HEALTH ADVISORY

Distributed via the CDC Health Alert Network
March 18, 2024, 12:30 PM ET
CDCHAN-00504

**Increase in Global and Domestic Measles Cases and Outbreaks:
Ensure Children in the United States and Those Traveling
Internationally 6 Months and Older are Current on MMR Vaccination**

Summary

The Centers for Disease Control and Prevention (CDC) is issuing this Health Alert Network (HAN) Health Advisory to inform clinicians and public health officials of an increase in global and U.S. measles cases and to provide guidance on measles prevention for all international travelers aged ≥ 6 months and all children aged ≥ 12 months who do not plan to travel internationally. Measles (rubeola) is highly contagious; one person infected with measles can infect 9 out of 10 unvaccinated individuals with whom they come in close contact. From January 1 to March 14, 2024, CDC has been notified of 58 confirmed U.S. [cases of measles](#) across 17 jurisdictions, including seven outbreaks in seven jurisdictions compared to 58 total cases and four outbreaks reported the entire year in 2023. Among the 58 cases reported in 2024, 54 (93%) were linked to international travel. Most cases reported in 2024 have been among children aged 12 months and older who had not received measles-mumps-rubella (MMR) vaccine. Many countries, including travel destinations such as Austria, the Philippines, Romania, and the United Kingdom, are experiencing measles outbreaks. To prevent measles infection and reduce the risk of community transmission from importation, all U.S. residents traveling internationally, regardless of destination, should be current on their MMR vaccinations. Healthcare providers should ensure children are current on routine immunizations, including MMR. Given currently high population immunity against measles in most U.S. communities, the risk of widescale spread is low. However, pockets of low coverage leave some communities at higher risk for outbreaks.

Background

[Measles](#) is a highly contagious viral illness and can cause severe health complications, including pneumonia, encephalitis (inflammation of the brain), and death, especially in unvaccinated persons. Measles typically begins with a prodrome of fever, cough, coryza (runny nose), and conjunctivitis (pink eye), lasting 2 to 4 days before rash onset. The incubation period for measles from exposure to fever is usually about 10 days (range 7 to 12 days), while rash onset is typically visible around 14 days (range 7 to 21 days) after initial exposure. The virus is transmitted through direct contact with infectious droplets or by airborne spread when an infected person breathes, coughs, or sneezes, and can remain infectious in the air and on surfaces for up to 2 hours after an infected person leaves an area. Individuals infected with measles are contagious from 4 days before the rash starts through 4 days afterward.

Declines in measles vaccination rates globally have increased the risk of measles outbreaks worldwide, including in the United States. Measles cases continue to be brought into the United States by travelers who are infected while in other countries. As a result, domestic measles outbreaks have been reported in most years, even following the declaration of U.S. [measles elimination](#) in 2000. Most importations come from unvaccinated U.S. residents.

Measles is almost entirely preventable through vaccination. MMR vaccines are safe and highly effective, with two doses being 97% effective against measles (one dose is 93% effective). When more than 95% of people in a community are vaccinated (coverage $>95\%$) most people are protected through community immunity (herd immunity). However, [vaccination coverage among U.S. kindergartners](#) has decreased from 95.2% during the 2019–2020 school year to 93.1% in the 2022–2023 school year, leaving

approximately 250,000 kindergartners susceptible to measles each year over the last three years. Thirty-six states plus the District of Columbia (DC) had less than 95% MMR coverage among kindergartners during the 2022–2023 school year. Of states with less than 95% MMR coverage, ten reported more than 5% of kindergartners had medical and nonmedical exemptions, highlighting the importance of targeted efforts at increasing [vaccine confidence](#) and access.

Recommendations for Healthcare Providers

- Schools, early childhood education providers, and healthcare providers should work to ensure students are current with [MMR vaccine](#).
 - Children who are not traveling internationally should receive their first dose of MMR at age 12 to 15 months and their second dose at 4 to 6 years.
- All U.S. residents older than age 6 months without evidence of immunity who are planning to travel internationally should receive MMR vaccine prior to departure.
 - Infants aged 6 through 11 months should receive one dose of MMR vaccine before departure. Infants who receive a dose of MMR vaccine before their first birthday should receive two more doses of MMR vaccine, the first of which should be administered when the child is age 12 through 15 months and the second at least 28 days later.
 - Children aged 12 months or older should receive two doses of MMR vaccine, separated by at least 28 days.
 - Teenagers and adults without evidence of measles immunity should receive two doses of MMR vaccine separated by at least 28 days.
- At least one of the following is considered evidence of measles immunity for international travelers: 1) birth before 1957, 2) documented administration of two doses of live measles virus vaccine (MMR, MMRV, or other measles-containing vaccine), or 3) laboratory (serologic) proof of immunity or laboratory confirmation of disease.
- Consider measles as a diagnosis in anyone with fever ($\geq 101^{\circ}\text{F}$ or 38.3°C) and a generalized maculopapular rash with cough, coryza, or conjunctivitis who has recently been abroad, especially in countries with ongoing [outbreaks](#). When considering measles, then:
 - **Isolate:** Do not allow patients with suspected measles to remain in the waiting room or other common areas of a healthcare facility; isolate patients with suspected measles immediately, ideally in a single-patient airborne infection isolation room (AIIR) if available, or in a private room with a closed door until an AIIR is available. Healthcare providers should be adequately protected against measles and should adhere to standard and airborne precautions when evaluating suspect cases, regardless of their vaccination status. Healthcare providers without evidence of immunity should be excluded from work from day 5 after the first exposure until day 21 following their last exposure. Offer testing outside of facilities to avoid transmission in healthcare settings. Call ahead to ensure immediate isolation for patients referred to hospitals for a higher level of care.
 - **Notify:** Immediately notify state, tribal, local, or territorial health departments ([24-hour Epi On Call contact list](#)) about any suspected case of measles to ensure rapid testing and investigation. States report measles cases to CDC.
 - **Test:** Follow CDC's testing recommendations and collect either a nasopharyngeal swab, throat swab, and/or urine for reverse transcription polymerase chain reaction (RT-PCR) and a blood specimen for serology from all patients with clinical features compatible with measles. RT-PCR is available at many state public health laboratories, through the APHL Vaccine Preventable Disease Reference Centers, and at CDC. Given potential shortages in IgM test kits, providers should be vigilant in contacting their state or local health department for guidance on testing.
 - **Manage:** In coordination with local or state health departments, provide appropriate measles post-exposure prophylaxis (PEP) as soon as possible after exposure to close contacts without evidence of immunity, either with MMR (within 72 hours) or immunoglobulin (within 6 days). The [choice of PEP](#) is based on elapsed time from exposure or medical contraindications to vaccination.

Recommendations for Health Departments

Measles is an immediately notifiable disease. State, tribal, local, and territorial health departments have the lead in disease investigations and should report measles cases and outbreaks within 24 hours through the state health department to CDC (measlesreport@cdc.gov) and through [NNDSS](#).

- Establish measles case reporting from healthcare facilities, providers, and laboratories to public health authorities.
- If measles is identified, conduct active surveillance for additional (secondary) cases and facilitate transportation of specimens immediately to confirm diagnosis.
- Record and report details about cases of measles, including adherence to recommended precautions and facility location(s) of index and secondary cases.
- Enhance outreach and communications to under-vaccinated communities through trusted messengers.

Recommendations for Parents and International Travelers

- Even if not traveling, ensure that children receive all recommended doses of MMR vaccine. Two doses of MMR vaccine provide better protection (97%) against measles than one dose (93%). Getting MMR vaccine is much safer than getting measles, mumps, or rubella.
- Anyone who is not protected against measles is at risk of getting infected when they travel internationally. Before international travel, check your [destination](#) and CDC's [Global Measles Travel Health Notice](#) for more travel health advice, including where measles outbreaks have been reported.
- Parents traveling internationally with children should consult with their child's healthcare provider to ensure that they are current with their MMR vaccinations at least 2 weeks before travel. Infants aged 6 to 11 months should have one documented dose and children aged 12 months and older should have two documented doses of MMR vaccine before international travel. Depending on where you are going and what activities you plan, other vaccines may be recommended too.
- After international travel, watch for signs and symptoms of measles for 3 weeks after returning to the United States. If you or your child gets sick with a rash and a high fever, call your healthcare provider. Tell them you traveled to another country and whether you or your child have received MMR vaccine.

For More Information

- Parents and International Travelers
 - [Measles Vaccines for Children | CDC](#)
 - [Plan for Travel – Measles | CDC](#)
 - [Global Measles Situation | CDC](#)
- Health Departments and Public Health Professionals
 - [Measles: Information for Public Health Professionals | CDC](#)
 - [CDC Measles Toolkit for Health Departments](#)
 - [Partnering for Vaccine Equity | CDC](#)
 - [Vaccine Preventable Diseases | APHL](#)
- Healthcare Providers
 - [Measles One-Pager for Healthcare Providers | Project Firstline and AAP](#)
 - [Immunization Schedules | CDC](#)
 - [Safety Information for Measles, Mumps, Rubella \(MMR\) Vaccines | CDC](#)
 - [For Healthcare Professionals – Diagnosing and Treating Measles | CDC](#)
 - [Interim Measles Infection Prevention Recommendations in Healthcare Settings | CDC](#)
 - [Measles – Vaccine Preventable Diseases Surveillance Manual | CDC](#)
 - [Rubeola / Measles | CDC Yellow Book 2024](#)
 - [Measles Lab Tools | CDC](#)
 - [Measles Serology | CDC](#)
 - [Measles Specimen Collection, Storage, and Shipment | CDC](#)
 - [Test Directory | Submitting Specimens to CDC | Infectious Diseases Laboratories | CDC](#)

- [Webinar Thursday, August 17, 2023 - We Must Maintain Measles Elimination in the United States: Measles Clinical Presentation, Diagnosis, and Prevention \(cdc.gov\)](#) (Free CE)

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network Messages

Health Alert	Conveys the highest level of importance about a public health incident.
Health Advisory	Provides important information about a public health incident.
Health Update	Provides updated information about a public health incident.

Montana Public Health Laboratory Guidance: Measles Testing



*State communicable disease reporting rules require health care providers suspecting measles to report suspected cases to local public health authorities **immediately**. Public health authorities may implement immediate control measures to prevent transmission and/or arrange immediate transport of the specimen when warranted.*

Specimen Criteria

Collect the following specimens to test for measles infection:

- Respiratory Specimen (Throat, NP, Nasal Swab)
- Serum

Specimen Collection for PCR Testing:

Collect specimens as soon as possible after appearance of rash, and ideally within 3 days of rash onset. Detection can be possible up to day 7 following onset of rash. *Respiratory Specimen:* Throat, Nasopharyngeal, or Nasal Dacron swabs in viral transport media.

Consult with Public Health authorities regarding PCR testing prior to rash development of individuals who may have had a recent exposure to measles.

Specimen Collection for IgM Testing:

For IgM testing, specimens must be collected >48 hours post rash onset.

- *Serum:* 1 – 2 ml of serum. Can be sent in a spun serum separator tube or can be poured off into a transport tube.

Transport Conditions:

- Keep *Respiratory specimens* cold, and transport with cold packs as soon as possible following specimen collection. Avoid repeat freeze-thaw cycles. If specimen transport is going to be delayed >24 hours, freeze the sample at -70°C and ship on dry ice.
- *Serum* specimens can be shipped cold (refrigerated) or frozen (preferable for IgM testing).

Submission Reminders:

*Please be sure to include the collection date and at least two patient identifiers (Name and DOB or medical record #) on the sample container.

*Use the online portal to order found at <https://labportal.hhs.mt.gov>. *For respiratory specimens, order Measles PCR. *For serology specimens, order Rubella IgM Serology (ND). *Be sure to select the appropriate specimen source when ordering. *For the Onset Date, enter the date of rash onset.

*Print the Requisition form, verify patient identifiers match the sample, and place the paperwork in the side pouch of the specimen bag. *Place the two specimens in separate specimen bags with the correct paperwork in each pouch. *Finally, create a manifest for all samples being shipped and place the manifest separately in the shipping container.

Specimens can be transported by courier (if available), UPS or FedEx to:

Montana Public Health Laboratory
1400 Broadway, Room B126
Helena, MT 59601

Please contact the Montana Public Health Laboratory at 1-800-821-7284 for more information and remember to report any suspect measles to your local health department.