

DPHHS HAN

ADVISORY

Cover Sheet



DATE

April 23, 2024

SUBJECT

Adverse Effects Linked to Counterfeit or Mishandled Botulinum Toxin Injections

INSTRUCTIONS

DISTRIBUTE to your local HAN contacts. This HAN is intended for general sharing of information.

- Time for Forwarding: **As Soon As Possible**
- Please forward to DPHHS at hhshan@mt.gov
- Remove this cover sheet before redistributing and replace it with your own

For LOCAL HEALTH
DEPARTMENT reference only
DPHHS Subject Matter Resource for
more information regarding this HAN,
contact:

DPHHS CDCP

Epidemiology Section
1-406-444-0273

*For technical issues related to the HAN
message contact the Emergency
Preparedness Section
at 1-406-444-0919*

DPHHS HAN Website:
<https://dphhs.mt.gov/publichealth/han>

REMOVE THIS COVER SHEET BEFORE REDISTRIBUTING AND REPLACE IT WITH YOUR OWN

Please ensure that DPHHS is included on your HAN distribution list.
hhshan@mt.gov

Categories of Health Alert Messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.

Information Service: passes along low level priority messages that do not fit other HAN categories and are for informational purposes only.

Please update your HAN contact information on the Montana Public Health Directory

DPHHS HAN

Information Sheet



DATE

April 23, 2024

SUBJECT

Adverse Effects Linked to Counterfeit or Mishandled Botulinum Toxin Injections

BACKGROUND

See CDC HAN 00507

INFORMATION

Twenty-two people in 11 U.S. states have reported adverse health effects after receiving injections with counterfeit botulinum toxin, or injections administered by unlicensed or untrained individuals, or in non-healthcare settings such as homes and spas. There may be symptom overlap between the adverse health effects from counterfeit or mishandled botulinum toxin injections and the early symptoms of botulism. None of the 22 cases that are part of this investigation were diagnosed in Montana residents.

RECOMMENDATIONS

Recommendations for Clinicians

Read CDC HAN 00507

Clinicians should be aware of symptom overlap between the presentation of localized adverse effects from injection of botulinum toxin and the early symptoms of botulism described in CDC HAN 00507. A suspected case of botulism is a clinical and public health emergency. Immediately call your [local or tribal public health department](#) with any suspect botulism cases, or with any instances of adverse health effects after administration of botulinum toxin.

Recommendations for Local and Tribal Health Departments

Read CDC HAN 00507

If a clinician suspects botulism, or other adverse health effects after administration of botulinum toxin, notify the Montana Communicable Disease Epidemiology Section, 24/7, at 406-444-0273.

This is an official
CDC HEALTH ADVISORY

Distributed via the CDC Health Alert Network
April 23, 2024, 11:00 AM ET
CDCHAN-00507

Adverse Effects Linked to Counterfeit or Mishandled Botulinum Toxin Injections

Summary

The Centers for Disease Control and Prevention (CDC) is issuing this Health Alert Network (HAN) Health Advisory to alert clinicians about risks of counterfeit or mishandled botulinum toxin injections. CDC, the U.S. Food and Drug Administration (FDA), and state and local partners are investigating clusters of 22 people in 11 U.S. states reporting adverse effects after receiving injections with counterfeit botulinum toxin or injections administered by unlicensed or untrained individuals or in non-healthcare settings, such as homes or spas. Eleven patients were hospitalized, and none have died. When botulinum toxin diffuses around the injection site, it can result in adverse effects. Botulism is the disease caused by botulinum toxin circulating in the blood and producing effects remotely from the injection site. There may be symptom overlap between the presentation of localized adverse effects from injection of botulinum toxin, especially in the head and neck, and the early symptoms of botulism. Information about the botulinum toxin injection (e.g., dose) can help distinguish between botulism and localized adverse effects but is challenging to obtain for counterfeit products. Clinicians and health departments should consider the possibility of adverse effects from botulinum toxin injections in patients presenting with localized paralysis. Clinicians should immediately contact their state, tribal, local, or territorial health department if they suspect botulism.

Background

Botulism is a rare and sometimes fatal illness caused by botulinum toxin. Initial botulism symptoms may include double or blurred vision, drooping eyelids, slurred speech, difficulty swallowing, and difficulty breathing. These symptoms may be followed by a descending, symmetric muscle weakness that progresses over hours to days. Administration of botulism antitoxin can neutralize toxin circulating in the blood; therefore, treating botulism patients with botulism antitoxin early in the course of disease can prevent the progression of paralysis and consequent complications. Administration of antitoxin is not indicated for local effects of low-dose injections of botulinum toxin preparations, because the low doses of injected toxin are not likely to reach circulation or produce botulism with its life-threatening manifestations.

Some localized paralytic effects, resulting from diffusion of the toxin around the injection site, are expected from botulinum toxin administration. Most individuals with localized symptoms (e.g., dysphagia after injection to the neck) following low-dose cosmetic or therapeutic injections using FDA-approved products will not require treatment with botulism antitoxin. Iatrogenic botulism can occur after cosmetic or therapeutic injections of botulinum toxin when the toxin circulates in the blood and produces effects remotely from the injection site. Iatrogenic botulism is rare; the most recent laboratory-confirmed domestic case occurred in 2017.

As of April 18, 2024, 22 people with adverse effects have been reported in California, Colorado, Florida, Illinois, Kentucky, Nebraska, New Jersey, New York City, Tennessee, Texas, and Washington. Symptom onset dates ranged from November 4, 2023, to March 31, 2024. All symptomatic people were females aged 25 to 59 years. All reported receiving botulinum toxin injections by unlicensed or untrained individuals or in non-healthcare settings, including homes or spas. Most (20, 91%) reported receiving botulinum toxin injections for cosmetic purposes.

Among all 22 people, symptoms began a median of 3 days after exposure (range 0 to 20 days) and included symptoms near the injection site (e.g., blurred vision and ptosis after facial injection), dry mouth, slurred speech, shortness of breath, fatigue, and generalized weakness. Of 20 people with information available, 11 (55%) symptomatic people were hospitalized. Six of the 22 symptomatic people received botulism antitoxin to treat suspected botulism. Seven symptomatic people underwent botulinum toxin testing to determine if they had circulating botulinum toxin, which would support a diagnosis of botulism; results were negative for six symptomatic people and are pending for one symptomatic person. Negative results do not rule out botulism as levels of toxin in serum may have fallen below the limit of detection of laboratory tests. None of the 22 symptomatic people met the [case definition for botulism](#), and none have died.

These adverse events have been linked to improper procurement and administration of botulinum toxin. Botulinum toxin should be administered only by licensed providers, using only recommended doses of FDA-approved botulinum toxin, preferably in a licensed or accredited healthcare setting. Providers should be trained in the proper administration of botulinum toxin, practicing in accordance with state and local requirements.

More information about the counterfeit products [may be found on FDA's website](#).

Recommendations for Clinicians

Diagnosis, consultation, and treatment

- Consider the possibility of adverse effects from botulinum toxin injections, including those given for cosmetic reasons, in patients presenting with localized paralysis near the injection site.
 - Ask patients about history of botulinum toxin injections, including the dose.
- Be aware of symptom overlap between the presentation of localized adverse effects from injection of botulinum toxin and the early symptoms of botulism. To help distinguish early botulism symptoms from localized adverse effects:
 - Assess for symmetry of cranial nerve palsies; symmetric cranial nerve palsies are expected with botulism.
 - Assess for progression of cranial nerve palsies, possibly followed by a descending symmetric flaccid paralysis. These should raise suspicion for botulism.
- **If botulism is suspected, call your [health department](#) immediately** for consultation. Health departments and health care providers can contact the CDC clinical botulism service 24/7 at 770-488-7100.
- If public health clinical consultation supports botulism, request antitoxin and begin treatment as soon as it is available. **Do not wait for laboratory confirmation to begin treatment.**

Clinician reporting

- A suspected case of botulism is a clinical and public health emergency. Suspected botulism cases should be reported immediately to the [24-hour emergency contact of your health department](#).
- Report adverse events related to the use of any medications, including suspected counterfeit medications, to FDA's [MedWatch Safety Information and Adverse Event Reporting Program](#).

Counseling patients

- Counsel patients who report using or being interested in using botulinum toxin about potential adverse effects.
- Advise patients to receive injections only from licensed providers who are trained in proper administration of FDA-approved botulinum toxin products, preferably in a licensed or accredited healthcare setting.

Recommendations for Laboratories

- Diagnostic testing for suspected botulism may be done through the [CDC National Botulism Laboratory](#) or state public health laboratories.
- Laboratory confirmation of botulism is done by demonstrating the presence of botulinum toxin in serum through either mouse bioassay or mass spectrometry.
- Testing varies by state. Contact the CDC clinical botulism service (available 24/7 at 770-488-7100) or your state health department for further guidance on submitting clinical specimens for testing.

Recommendations for Public Health Professionals

- Health departments should contact the CDC clinical botulism service 24/7 at 770-488-7100 for consultation and antitoxin release.
- Health departments should report botulism cases to CDC through the [National Notifiable Diseases Surveillance System \(NNDSS\)](#).
- Contact botsurveillance@cdc.gov for information on how to report adverse effects following administration of unverified or counterfeit botulinum toxin products, by unlicensed or untrained individuals, or in non-healthcare settings such as homes or spas.

Recommendations for the Public

- Get injections only from licensed and trained professionals in licensed or accredited healthcare settings.
- If you are concerned that you or someone you know might have [symptoms of botulism](#), including trouble swallowing or breathing, see your doctor or go **immediately** to the emergency room. **Do not wait.**
- Report suspected counterfeit botulinum toxin products to FDA at 800-551-3989 or through FDA's [form for reporting suspected criminal activity](#).
- Report harmful reactions related to the use of any medications, including suspected counterfeit medications, to FDA's [MedWatch Safety Information and Adverse Event Reporting Program](#).

For More Information

- [CDC: Harmful Reactions Linked to Counterfeit “Botox” or Mishandled Botulinum Toxin Injections](#)
- [FDA: Counterfeit Version of Botox Found in Multiple States](#)
- [Information for Health Professionals on Botulism | CDC](#)
- [About Botulism | CDC](#)
- [Preventing Botulism | CDC](#)
- [Injection Safety | CDC](#)

References

1. Halai U, Terashita D, Kim M, et al. Notes from the Field: Intestinal Colonization and Possible Iatrogenic Botulism in Mouse Bioassay-Negative Serum Specimens — Los Angeles County, California, November 2017. *MMWR Morb Mortal Wkly Rep*. 2018;67(43):1221–1222. doi:[10.15585/mmwr.mm6743a6](#)
2. Rao AK, Sobel J, Chatham-Stephens K, Luquez C. Clinical Guidelines for Diagnosis and Treatment of Botulism, 2021. *MMWR Recomm Rep*. 2021;70(No. RR-2):1–30. doi:[10.15585/mmwr.rr7002a1](#)
3. Rao AK, Lin NH, Jackson KA, Mody RK, Griffin PM. Clinical Characteristics and Ancillary Test Results Among Patients With Botulism — United States, 2002–2015. *Clin Infect Dis*. 2018;66(suppl_1):S4–S10. doi:[10.1093/cid/cix935](#)

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on

critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages

Health Alert	Conveys the highest level of importance about a public health incident.
Health Advisory	Provides important information about a public health incident.
Health Update	Provides updated information about a public health incident.